# A Toronto Midwifery Practice's Postpartum Visit Schedule and Chest/ Breastfeeding Rates before and during the COVID-19 Pandemic: A Retrospective Cohort Study

Claire Osepchook<sup>1</sup>, Wissutta Durongphongtorn<sup>2</sup>, Claire Dion Fletcher<sup>3</sup>

<sup>1</sup>Registered Midwife, Seventh Generation Midwives Toronto; <sup>2</sup>Registered Midwife; <sup>3</sup>Registered Midwife, Seventh Generation Midwives Toronto and Assistant Professor in the Midwifery Education Program at the Toronto Metropolitan University

\*Corresponding author: Claire Osepchook: cosepchook@sgmt.ca

Submitted: 2 May 2025; Accepted: 9 October 2025; Published: 1 November 2025

#### **ABSTRACT**

**Objective:** This study aims to determine whether the alternate midwifery postpartum visit schedule imposed by the COVID-19 pandemic affected the chest/breastfeeding rate of midwifery clients compared to the standard postpartum visit schedule.

**Methods:** Three hundred anonymized records of midwifery clients at a Toronto midwifery practice were analyzed for the total number of postpartum parent-infant visits and method of infant feeding at Day 3, 2 weeks, and 6 weeks postpartum. The pre-pandemic cohort  $(n_1 = 150)$  gave birth from January to October 2018. The pandemic cohort  $(n_2 = 150)$  gave birth from April to November 2020.

**Results:** The cohort comparison showed that the average number of postpartum visits during the COVID-19 pandemic remained unchanged. The exclusive chest/breastfeeding rate at 6 weeks postpartum was not significantly different, and it remained high.

**Conclusion:** Midwifery clients and midwives persevered through the COVID-19 pandemic, demonstrating a high commitment to protecting individualized dyad-centred care and in-person chest/breastfeeding support.

#### RÉSUMÉ

**Objectif :** Cette étude vise à déterminer si le calendrier alternatif des visites postnatales imposé par la pandémie de COVID-19 a eu une incidence sur le taux d'allaitement maternel des clientes des sages-femmes par rapport au calendrier standard des visites postnatales.

**Méthodes :** Trois cents dossiers anonymisés de clientes d'une pratique de sages-femmes à Toronto ont été analysés afin de déterminer le nombre total de visites postnatales des parents et des nourrissons et

la méthode d'alimentation des nourrissons au troisième jour, à deux semaines et à six semaines après l'accouchement. La cohorte pré-pandémique (n1 = 150) a accouché entre janvier et octobre 2018. La cohorte pandémique (n2 = 150) a accouché entre avril et novembre 2020.

**Résultats :** La comparaison des cohortes a montré que le nombre moyen de visites post-partum pendant la pandémie de COVID-19 est resté inchangé. Le taux d'allaitement maternel exclusif à 6 semaines après l'accouchement n'était pas significativement différent et est resté élevé.

**Conclusion :** Les clientes des sages-femmes et les sages-femmes ont persévéré pendant la pandémie de COVID-19, démontrant un engagement fort en faveur de la protection des soins individualisés centrés sur la dyade et du soutien à l'allaitement maternel en personne.

#### INTRODUCTION

#### The COVID-19 Pandemic

On March 11, 2020, the World Health Organization (WHO) declared a global pandemic because of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), commonly known as COVID-19.1 The world had to rapidly adapt to the constant change in restrictions and policies to limit the transmission of the virus.<sup>1</sup> In Ontario, the provincial government implemented public health measures such as gathering restrictions, physical distancing, workfrom-home policies, suspension and delay of many health care services, increase in virtual health care, and mandatory use of face masks in indoor public spaces.<sup>2</sup> COVID-19 had widespread health effects across communities. Public Health Ontario epidemiological reports found that marginalized communities in the most materially deprived neighbourhoods and the most ethnically diverse neighbourhoods in Ontario experienced higher rates of COVID-19 infection, more severe outcomes, and higher mortality rates.3,4

# Chest/breastfeeding rates during the COVID-19 Pandemic

Chest/breastfeeding has many well-known health benefits. The WHO, Health Canada, and the Canadian Paediatric Society recommend exclusive chest/breastfeeding for the first 6 months with continued chest/breastfeeding for up to 2 years and beyond. 5-7 The benefits of chest/breastfeeding for the newborn include protection against gastrointestinal and respiratory infections, protection against sudden infant death syndrome, and enhanced cognitive development. 6-10 Chest/breastfeeding protects against breast cancer, ovarian cancer, and type 2 diabetes for the chest/breastfeeding parent. 6.8.10

In June 2020, the WHO recommended that chest/breastfeeding be initiated and continued in spite of a suspected or confirmed COVID-19 infection, as the risk of transmission of the virus to the infant was found to be low.<sup>11</sup> Newer research has demonstrated that chest/breastfeeding provides newborns with immunity against COVID-19 because of the presence of antibodies found in human milk.<sup>12</sup>

In spite of the COVID-19 pandemic, chest/ breastfeeding intentions during pregnancy remained high around the world.13 Yet, the impacts of the COVID-19 restrictions on chest/breastfeeding rates are mixed. In a United Kingdom sample, chest/ breastfeeding rates remained comparable to prepandemic rates.<sup>14</sup> Similarly, an international crosssectional study found that COVID-19-imposed changes in perinatal care did not affect chest/ breastfeeding rates.15 However, the same study found that virtual support groups were positively associated with continued chest/breastfeeding at 6 months, particularly for parents with their first child.15 They also found that a lack of support negatively affected chest/breastfeeding rates.<sup>15</sup> found that the COVID-19 pandemic public health restrictions mostly had negative impacts on chest/ breastfeeding such as separation of parent and infant, lack of skin to skin contact, and insufficient postpartum support.16 Another study found the most common reason for chest/breastfeeding cessation during the COVID-19 pandemic was the lack of professional face-to-face support.<sup>17</sup>

There were also some unexpected positive impacts of the pandemic restrictions mainly reported by participants with higher socioeconomic status, which included protecting the chest/breastfeeding parent-infant dyad from unwanted visitors at home, more time to focus on chest/breastfeeding,

and greater support from their partner with delayed return to work outside the home.<sup>16</sup>

# The Standard Ontario Midwifery Postpartum Care Visit Schedule

The Ontario midwifery model of postpartum care provides high-quality primary care for the parentinfant dyad from birth to 6 weeks postpartum. The hallmarks of this model are multiple visits, visits in the client's preferred setting (home, hospital, or clinic), and individualized dyad-centered care.<sup>18</sup> Midwives provide clinical care for the parent and newborn, and support mental health well-being, infant growth and development, chest/breastfeeding, and parent-infant bonding. A typical postpartum visit schedule includes three to five visits in home or hospital and one to three clinic visits. <sup>18</sup> In 2016–2017, midwives in Ontario provided an average of 6.6 postpartum visits per client with a range from 0-25 visits.18 The College of Midwives of Ontario rescinded the 2015 Postpartum/ newborn Visits Standard<sup>19</sup> in 2018, to better support midwives to use their clinical judgment and adjust the postpartum visit schedule to respond to the needs of clients.<sup>18</sup> The Ontario midwifery postpartum care model has resulted in excellent health outcomes for both parents and infants, including high chest/ breastfeeding rates at 6 weeks postpartum,18 high client satisfaction,18 and reduced unscheduled postpartum hospital emergency department use.20

# Alternative Ontario Midwifery Postpartum Care Visit Schedule

Beginning in March 2020, the Association of Ontario Midwives (AOM) proposed an alternative midwifery postpartum visit schedule to reduce COVID-19 exposure and community transmission using guidance from the WHO.<sup>21</sup> The alternative schedule proposed one in-person visit within 48 hours of birth, at least one additional in-person visit in the first week, and additional virtual visits, including the discharge visit, by phone or videoconference.<sup>21</sup> Health precautions were recommended for clinical situations that required an in-person assessment by the midwife.

# **OBJECTIVE**

This study aims to determine whether the alternative midwifery postpartum visit schedule imposed

by the COVID-19 pandemic affected the chest/ breastfeeding rate of midwifery clients compared to the *standard postpartum visit schedule*.

# **METHOD**

## Setting

Seventh Generation Midwives Toronto (SGMT) is an Indigenous-led midwifery practice group of Indigenous registered midwives, Indigenous midwives working under an exemption of the Midwifery Act, and non-Indigenous registered midwives located in Toronto, Ontario.

#### Design

## Literature Review

A literature review was conducted from August 2022 to 2024 using the following databases: Cochrane Library, PubMed MEDLINE, MEDLINE, Cumulative Index to Nursing & Allied Health (CINAHL) EBSCO, the Toronto Metropolitan University Libraries, and Google Scholar. The main search terms included: "midwi\* AND postpartum AND (visit OR schedule) AND (breastfeed\* OR infant feed\* OR lact\*] AND COVID," with a preference for relevant articles in Ontario, Canada. The database search limits were English language and peerreviewed articles. To our knowledge, this study is the first to research the Ontario midwifery postpartum visit schedule and its effect on breastfeeding rates during the COVID-19 pandemic. The literature review found no relevant studies.

### Study Design

This is a quantitative retrospective cohort study of midwifery care for clients who gave birth before or during the COVID-19 pandemic and had 6 weeks of postpartum care with SGMT midwives. The primary outcome is chest/breastfeeding status at 6 weeks postpartum, described as exclusive chest/breastfeeding, combined feeding [chest/breastfeeding and formula feeding], or exclusive formula feeding. The secondary outcomes include [1] the mean number of postpartum visits, [2] the type of postpartum visits, [3] feeding status at Day 3 postpartum, [4] feeding status at 2 weeks postpartum, and [5] the percentage of midwifery clients accessing additional chest/breastfeeding support.

#### **Participants**

**Exclusion Criteria** 

Midwifery clients with any of the following conditions were excluded from the study: [1] stillbirth, [2] preterm birth less than 37+0 weeks gestation, [3] neonatal intensive care [NICU] admission, [4] severe postpartum hemorrhage [>1000 mL], [5] hypertensive disorders of pregnancy, [6] preexisting or gestational diabetes, [7] maternal hospital readmission in the postpartum period, or [8] twin pregnancies.

# Sample Size

Using purposive sampling, 300 anonymized client charts, 150 in each cohort, were included to detect a statistically significant change of 10% in the chest/breastfeeding rate for a type-I error of 5% and a power of 90%.

#### **Data Collection**

A total of 300 midwifery clients (N = 300) passed the exclusion criteria, and data from their Ontario perinatal records and postnatal records were collected and anonymized. The pre-pandemic cohort ( $n_1$  = 150) gave birth between January 1, 2018 and October 31, 2018. The pandemic cohort ( $n_2$  = 150) gave birth between April 1, 2020 and November 30, 2020.

#### **Data Analysis**

The baseline characteristics and study outcomes were compared between the two cohorts to identify confounding variables. potential Continuous variables were compared using the independent t-test. Variance in continuous variables was compared using the F-test. Categorical variables were compared using the Chi-square test. Multivariable logistic regression analysis was used to determine whether the type of postpartum schedule was associated with any chest/ breastfeeding rate differences while adjusting for potential confounding factors. The results are presented as an adjusted odds ratio (aOR) with a 95% confidence interval (95% CI).

# **Ethics Approval**

The DAN Women & Babies Program Research Committee and the Sunnybrook Research Ethics

Board reviewed and approved this study, project identification #5409.

#### **RESULTS**

# **Demographics**

The pre-pandemic cohort consisted of 150 midwifery clients who gave birth between January and October 2018, while the pandemic cohort consisted of 150 clients who gave birth between April and November 2020. The demographics of the two cohorts are shown below in Table 1. There were no significant differences in age and ethnicity between the cohorts. The pandemic cohort had 16.0% more primiparous clients and 16.7% more clients without chest/breastfeeding experience than the pre-pandemic cohort. Since the primiparous clients had no chest/breastfeeding experience, the higher percentage of primiparous clients without chest/breastfeeding experience in the pandemic cohort was the main significant difference between the cohorts. The pandemic cohort also had a statistically significant 9.3% increase in home births.

#### **Chest/Breastfeeding Rates**

The exclusive chest/breastfeeding rate at 6 weeks postpartum was 82.7% in the pre-pandemic cohort and 74.7% in the pandemic cohort (Figure 1); an 8.0% decline during the pandemic but not statistically significant. The pandemic cohort had an 8.0% increase in combination feeding at 6 weeks postpartum compared to the pre-pandemic cohort. The exclusive formula feeding rate at 6 weeks postpartum was the same for both cohorts at 2.7%. There were no statistically significant differences between cohorts for exclusive chest/breastfeeding, combination feeding, and formula feeding rates at Day 3 and 2 weeks postpartum (Figure 1).

#### Midwifery Postpartum Visit Schedule

In the pre-pandemic cohort, midwives provided an average of six postpartum visits with a standard deviation of  $\pm$  1 similar to the standard midwifery postpartum visit schedule. The range of the number of postpartum visits provided was four to eight visits [Table 2]. During the pandemic, the average number of total postpartum visits provided by the midwives

**Table 1.** Demographic characteristics of the pre-pandemic cohort and the pandemic cohort.

Characteristic	Pre-pandemic cohort (n <sub>1</sub> = 150) Frequency (%)	Pandemic cohort (n <sub>2</sub> = 150) Frequency (%)	p-value $[\alpha < 0.05]$	
Age at delivery, mean (years ± SD)	32 ± 4.960	33 ± 4.444	0.067	
Age at delivery, range (years)	19 to 45	18 to 47	0.091	
Ethnicity				
White/European	70 [46.7%]	76 (50.7%)		
First Nations/Métis/Inuit/Mixed	22 (14.7%)	12 [8.0%]		
East/Southeast Asian	17 (11.3%)	16 (10.7%)		
Black/Caribbean	9 (6.0%)	8 (5.3%)		
Latin/Central/South American	7 (4.7%)	12 [8.0%]		
South Asian	6 [4.0%]	6 (4.0%)		
West Asian	2 [1.3%]	6 (4.0%)		
Multiple Ethnicities	17 (11.3%)	14 [9.3%]		
Parity*				
Primiparous	58 (38.7%)	82 [54.7%]		
Multiparous	92 (61.3%)	68 [45.3%]		
Chest/breastfeeding experience*				
No chest/breastfeeding experience	59 (39.3%)	84 [56.0%]		
Previous chest/breastfeeding experience	91 (60.7%)	66 [44.0%]		
Birth location*				
Sunnybrook Health Sciences Centre	115 (76.7%)	104 [69.3%]	0.153	
Toronto Birth Centre	29 (19.3%)	26 [17.3%]	0.654	
Home*	6 [4.0%]	20 (13.3%)	0.004	


SD, standard deviation; \*, statistically significant difference p < 0.05.

remained the same as before the pandemic, but there was a greater variation in the range of total number of postpartum visits from three to nine visits (Table 2).

In the pre-pandemic cohort, over 99% of the postpartum visits were in-person, while virtual visits made up less than 1% of the total (Table 3). Similarly, in the pandemic cohort, 99% of the postpartum visits were in-person while less than 1% were virtual (Table 3).

# Additional Chest/Breastfeeding Support

In the pandemic cohort, there was a statistically significant 10.7% increase in clients who accessed additional chest/breastfeeding support from lactation consultants or pediatricians (Figure 2). In addition, we found there was a statistically significant increase in the number of clients with no prior chest/breastfeeding experience in the pandemic group compared with the pre-pandemic group; 56% and 39.3%, respectively (Table 1).



**Figure 1.** Chest/breastfeeding rates at Day 3, 2 weeks, and 6 weeks postpartum of the pre-pandemic cohort and pandemic cohort.

Table 2. Total number of postpartum visits before and during the COVID-19 pandemic.

Measurement	Pre-pandemic cohort (n <sub>1</sub> = 150)	Pandemic cohort (n <sub>2</sub> = 150)
Total number of postpartum visits, mean ± SD	6 ± 0.872	6 ± 1.197
Total number of postpartum visits, range	4 to 8	3 to 9

SD, standard deviation; p-value = 1.000.

**Table 3.** Percentage of in-person and virtual postpartum visits before and during the COVID-19 pandemic.

Measurement	Pre-pandemic cohort	Pandemic cohort
Percentage of in-person postpartum visits	99.89%	99.4%
Percentage of virtual postpartum visits	0.11%	0.6%


p-value = 1.000.

#### DISCUSSION

We did not find a statistically significant difference in exclusive chest/breastfeeding rates at 6 weeks postpartum in the pre-pandemic and pandemic cohorts in this study. The rate of exclusive chest/breastfeeding in the pre-pandemic group was 82.7%, while the exclusive chest/breastfeeding rate in the pandemic group was 74.7%. These numbers are slightly higher than the 2022 Ontario midwifery rate for exclusive chest/breastfeeding at 6 weeks

[71%]<sup>22</sup> and higher than provincial and national rates of exclusive breastfeeding at 2 months, 68.4% and 66.6%, respectively.<sup>23,24</sup>

One of the hallmarks of the Ontario midwifery postpartum care model is the multiple in-person postpartum visits in the setting of the client's choice. Similar to the AOM's Postpartum Visit Schedule Guideline<sup>18</sup>; a large 2022 Cochrane Review of 116 trials investigating effective chest/breastfeeding support found that four to eight visits for chest/



**Figure 2.** Comparison of additional chest/breastfeeding support accessed among the pre-pandemic and pandemic cohorts.

breastfeeding support were associated with higher exclusive chest/breastfeeding at 4 to 6 weeks and 6 months.<sup>25</sup> In this study, the midwives provided between four to eight in-person postpartum visits, averaging six postpartum visits before the pandemic (Table 2), which was in line with Ontario midwifery guidelines at that time.

The AOM's "COVID-19: Reduced schedule of visits and use of PPE in midwifery contacts" guideline proposed two visits in the first week with additional visits by phone or videoconference.<sup>21</sup> Interestingly, we found that the average number of postpartum visits in the pandemic cohort was unchanged, in spite of the recommendation to decrease the number of visits to reduce COVID-19 exposure and community transmission. Similarly, the type of visit did not change between the cohorts, and it was almost exclusively in-person in spite of the recommendation for virtual visits when appropriate. The 2022 Cochrane Review mentioned above found that the type of support: face to face, telephone, digital technologies, or a combination, did not significantly affect chest/breastfeeding rates.<sup>25</sup> However, a study from the United Kingdom reported that 70.3% of participants attributed their decision for early chest/breastfeeding cessation during the pandemic to a lack of face-to-face chest/ breastfeeding support.<sup>17</sup> Parents in Canada who were able to access virtual chest/breastfeeding support during the COVID-19 pandemic unanimously viewed it as unhelpful.26

Midwives provide a full complement of postpartum care for both the birthing parent and infant. This unique model of care includes, but is not limited to, physical assessments such as blood pressure measurements and other vitals; perineal tear and c-section incision healing; initiation, establishment, and continuation of chest/breastfeeding support, including supplemental or formula feeding; newborn assessments, screens, medications, and prophylaxis; mental health support; and parent-infant bonding support. Many of these aspects of care would have been more difficult to provide using the alternative reduced visit schedule.

Other Ontario perinatal care providers have reported difficulties in providing standard clinical care virtually, such as the inability to perform physical assessments or hands-on emotional support.<sup>26</sup> They asserted that virtual care could not substitute in-person clinical care.<sup>26</sup>

The midwives in this study provided frequent inperson postpartum visits to effectively assess and support clients' and newborns' well-being in spite of the recommended reduced visit schedule during the COVID-19 pandemic. Midwives are supported by the CMO and AOM to adjust the postpartum visit schedule using clinical judgment and appropriate health precautions. A recent study examining midwifery care in Ontario during the COVID-19 pandemic showed that midwives adapted protocols and quidelines to uphold the tenets of midwifery

care during a crisis.<sup>27</sup> Similarly, the midwives in this study remained committed to meeting the needs of clients during the COVID-19 pandemic and provided frequent in-person postpartum care visits while ensuring that health precautions were in place such as wearing full personal protective equipment, implementing health screening strategies for clients, and performing hand hygiene and appropriate cleaning of clinical equipment.

In spite of the number of postpartum visits provided remaining the same in both cohorts, we found a 10.6% statistically significant increase in additional non-midwifery chest/breastfeeding support accessed by clients in the pandemic group. Most of the clients who accessed additional feeding support in both cohorts were primiparous parents without chest/breastfeeding experience. The additional support was provided by lactation consultants and pediatricians. The pandemic cohort had a 16% statistically significant higher number of primiparous clients compared to the pre-pandemic group which could explain the increase in support accessed outside of midwifery care.

Lastly, we found that compared to the prepandemic cohort, the pandemic cohort had a higher rate of home births. There was a 9.3% statistically significant increase in SGMT midwife-attended home births between April and November 2020. This was likely because of clients' desire to avoid COVID-19 exposure in clinical settings. Similarly, Ontario-wide data showed an increase in midwife-attended home births, peaking at 16.6% in May of 2020.28 Increases in out-of-hospital births may contribute to the initiation of and establishment of chest/breastfeeding because of reduction in interventions, increased skin-to-skin and decreased disruption of the parent-infant dyad.

## **CONCLUSION**

The high exclusive chest/breastfeeding rates of midwifery clients in Ontario have been attributed to the midwifery postpartum care model which provides individualized care for the parent-infant dyad. The COVID-19 pandemic resulted in many restrictions that affected the provision of midwifery care. Midwives in Ontario were advised

to decrease the number of postpartum visits as well as to replace in-person visits to virtual encounters when appropriate to reduce the exposure and transmission of the COVID-19 virus. Midwifery clients and midwives in this study demonstrated a high commitment to exclusive chest/breastfeeding before and during the pandemic. The pandemic cohort in this study was found to have similar exclusive chest/ breastfeeding rates as the pre-pandemic group. When we looked at the postpartum visits provided by the midwives, we found that the midwives did not reduce the number of visits nor the number of in-person visits for clients. Midwives are supported by the CMO to use clinical judgement and expertise in postpartum care when devising individualized postpartum care plans for clients. The midwives used a multiple in-person postpartum visits model to meet clients' needs, demonstrated by similar high exclusive chest/breastfeeding rates in the pre-pandemic and pandemic groups.

#### LIMITATIONS

This study's main methodological limitations are that the data were collected from one midwifery practice group in Toronto and during the first 10 months of a multiyear pandemic with constantly changing health recommendations and restrictions. The results may not be generalizable to other midwives in Canada because of differing models of midwifery postpartum care across the country. Future research on the Ontario midwifery postpartum care model, how it has been transformed by midwives during and since the COVID-19 pandemic, and how midwives determine the need for postpartum care would provide more insight into the excellent exclusive chest/breastfeeding rates and other health outcomes of midwifery clients in Ontario.

# **ACKNOWLEDGEMENTS**

We would like to thank Arel Ortiz Rosas, student midwife for her contributions to this work, as well as Dr. Nir Melamed and Tasha Macdonald, RM, for their guidance.

#### **CONFLICT OF INTEREST**

The authors declare no conflict of interest.

#### **FUNDING**

This research was funded by the DAN Women & Babies Program at Sunnybrook Health Sciences Centre in Toronto, Ontario.

#### **AUTHOR BIOGRAPHIES**

Claire Osepchook is a registered midwife practicing at Seventh Generation Midwives Toronto.

Claire Dion Fletcher is a registered midwife from Seventh Generation Midwives Toronto and an assistant professor in the Midwifery Education Program at the Toronto Metropolitan University. Claire sits on the Board of the Canadian Association of Midwives.

Wissutta Durongphongtorn is a registered midwife practicing at The Midwives Collective of Toronto. She was a student in the Midwifery Education Program at Toronto Metropolitan University when she contributed to this study.

#### **REFERENCES**

- World Health Organization. WHO Director-General's opening remarks at the media briefing on COVID-19 [Internet]. 2020 [cited 2024 Aug 15]. Available from: https://www.who.int/director-general/speeches/detail/ who-director-general-s-opening-remarks-at-the-mediabriefing-on-covid-19---11-march-2020
- Canadian Institute for Health Information. Canadian COVID-19 intervention timeline [Internet]. 2022 [cited 2024 Aug 16]. Available from: https://www.cihi.ca/en/ canadian-covid-19-intervention-timeline
- 3. Ontario Agency for Health Protection and Promotion (Public Health Ontario). COVID-19 in Ontario: A focus on neighbourhood material deprivation, February 26, 2020 to December 31, 2022 [Internet]. Toronto; 2023 May [cited 2024 Aug 15]. Available from: https://www.publichealthontario.ca/-/media/documents/ncov/epi/2020/06/covid-19-epi-material-deprivation.pdf?la=en
- 4. Ontario Agency for Health Protection and Promotion [Public Health Ontario]. COVID-19 in Ontario—a focus on neighbourhood diversity, February 26, 2020 to December 31, 2022 [Internet]. Toronto; 2023 May [cited 2024 Aug 15]. Available from: https://www.publichealthontario.ca/-/media/documents/ncov/epi/2020/06/covid-19-epi-diversity.pdf?la=en#:~:text=The%20case%20 fatality%20rate%20for,3.3%25%20[Table%202
- World Health Organization. Infant and young child feeding [Internet]. 2021 [cited 2022 Jul 6]. Available from: https://www.who.int/news-room/fact-sheets/detail/infant-and-young-child-feeding
- Infant Feeding Joint Working Group. Nutrition for healthy term infants: Recommendations from birth to six months [Internet]. 2022 [cited 2022 Aug 5]. Available from: https:// www.canada.ca/en/health-canada/services/canadafood-guide/resources/infant-feeding/nutrition-healthyterm-infants-recommendations-birth-six-months.html
- 7. Critch JN, Canadian Paediatric Society, Nutrition and Gastroenterology Committee . Nutrition for healthy term

- infants, birth to six months: An overview. Paediatr Child Health [Internet]. 2013 [cited 2022 Jul 21];18[4]: 206-7. https://doi.org/10.1093/pch/18.4.206
- Victora CG, Bahl R, Barros AJ, et al. Breastfeeding in the 21st century: Epidemiology, mechanisms, and lifelong effect. Lancet [Internet]. 2016 Jan 30 [cited 2022 Jul 24]; 387[10017]:475–90. https://doi.org/10.1016/S0140-6736 [15]01024-7
- Kramer MS, Kakuma R. Optimal duration of exclusive breastfeeding. Cochrane Database of Systematic Reviews [Internet]. 2012 Aug 15 [cited 2022 Jul 19];2012[8]:CD003517. https://doi.org/10.1002/14651858.CD003517.pub2
- Public Health Agency of Canada. Breastfeeding. In: Family-centered maternity and newborn care: National guidelines [Internet]. 2019 [cited 2024 Aug 21]; p.1–62. Available from: https://www.canada.ca/en/public-health/ services/publications/healthy-living/maternity-newborncare-guidelines-chapter-6.html
- World Health Organization. Breastfeeding and COVID-19 [Internet]. 2020 Jun [cited 2022 Jul 6]. Available from: https://www.who.int/publications/i/item/WHO-2019-nCoV-Sci\_Brief-breastfeeding-2020.1
- Richard RM, Maziashvili G, Tran M, et al. Breast milk conferred immunity to infants against COVID-19 [Internet]. 2023 Jul 18 [cited 2025 Aug 30];15[7]:e42075. https://doi. org/10.7759/cureus.42075
- Chien LY, Lee EY, Coca KP, et al. Impact of COVID-19 on breastfeeding intention and behaviour among postpartum women in five countries. Aust. J. Midwifery [Internet]. 2022 Jun 17 [cited 2022 Aug 11]; Available from: https://www.sciencedirect.com/science/article/pii/ S187151922200107X?via%3Dihub
- Turner S, McGann B, Brockway M. A review of the disruption of breastfeeding supports in response to the COVID-19 pandemic in five Western countries and applications for clinical practice. Int. Breastfeed J. [Internet]. 2022 Dec 1 [cited 2024 Aug 20];17[1]:1–13. https://doi.org/10.1186/ s13006-022-00478-5
- Ganho-Ávila A, Guiomar R, Sobral M, et al. The impact of COVID-19 on breastfeeding rates: An international cross-sectional study. Midwifery [Internet]. 2023 May 1 [cited Jan 2025 30];120:103631. https://doi.org/10.1016/j. midw.2023.103631
- Lubbe W, Niela-Vilén H, Thomson G, et al. Impact of the COVID-19 pandemic on breastfeeding support services and women's experiences of breastfeeding: A review. Int. J. Women's Health [Internet]. 2022 Oct [cited 2024 Aug 16];14:1447–57. https://doi.org/10.2147/IJWH.S342754
- Brown A, Shenker N. Experiences of breastfeeding during COVID 19: Lessons for future practical and emotional support. Matern. Child Nutr. [Internet]. 2021 Jan 1 [cited 2022 Jul 27];17[1]:e13088. https://doi.org/10.1111/mcn.13088
- Midwifery Task Force on Postpartum Visit Schedules. Postpartum visit schedules: Guideline for timing postpartum visits. Association of Ontario Midwives [Internet].
   2019 [cited 2022 Jul 21];1-11. Available from: https://www.ontariomidwives.ca/sites/default/files/2019-07/Postpartum%20visit%20schedules%20web\_V06.pdf
- College of Midwives of Ontario. Postpartum/newborn visits [Internet]. 2015 [cited 2022 Jul 21]. Available from: https://www.cmo.on.ca/wp-content/uploads/2015/07/ Final-PP-Newborn-visits-Standard.pdf
- Sorbara C, Ray JG, Darling EK, et al. Postpartum emergency department use following midwifery-model vs obstetricsmodel care. JAMA. Netw. Open [Internet]. 2024 Apr 29 [cited 2025 Jan 30];7(4): e248676. https://doi.org/10.1001/ jamanetworkopen.2024.8676

- 21. Association of Ontario Midwives. COVID-19: Reduced schedule of visits and use of PPE in midwifery contacts [Internet]. 2020 Mar [cited 2024 Aug 15]. Available from: https://www.ontariomidwives.ca/sites/default/files/IPAC%20Protocols/2020%2003%2020%20COVID19-Reduced%20schedule%20of%20visits%20and%20use%20of%20PPE.pdf
- 22. Association of Ontario Midwives (AOM). Midwifery by the numbers [Internet]. Toronto: AOM; 2020 [cited 2021 Mar 16]. Available from: https://www.ontariomidwives.ca/midwifery-numbers
- 23. Health Nexus. Breastfeeding in Ontario, fact sheet #1: Notable trends within the province [Internet]. Toronto; 2015 [cited 2022 Jul 20]. Available from: https://resources.beststart.org/product/b15e-breastfeeding-ontario-factsheet1-notable-trends/
- 24. Government of Canada. Canada's breastfeeding dashboard 2024 edition [Internet]. 2024 Oct 31 [cited 2025 May 6]. Available from: https://health-infobase.canada.ca/breastfeeding/#a3

- Gavine A, Shinwell SC, Buchanan P, et al. Support for healthy breastfeeding mothers with healthy term babies. Cochrane Database of Systematic Reviews [Internet]. 2022 Oct 25 [cited 2024 Aug 20];2022[10]. https://doi. org/10.1002/14651858.CD001141.pub6
- 26. Rice K, Williams S. Women's postpartum experiences in Canada during the COVID-19 pandemic: A qualitative study. CAMJ Open [Internet]. 2021 Apr 1 [cited 2024 Aug 16];9(2): E556-62. https://doi.org/10.9778/cmajo.20210008
- 27. Macdonald M, Burton N, Aseffa F, et al. Equity in care: Midwifery in Ontario during the COVID-19 pandemic. CJMRP [Internet]. 2024 Aug 24 [cited 2025 Feb 5];23[1]: 89–103. https://doi.org/10.22374/cjmrp.v23i1.16
- 28. Association of Ontario Midwives. #Midwifery data matters: Examining the rise in out-of-hospital birth during the COVID-19 pandemic [Internet]. 2022 [cited 2024 Aug 20]. Available from: https://www.ontariomidwives.ca/midwiferydatamatters-examining-rise-out-hospital-birth-during-covid-19-pandemic